

The evaluation of stability and maturity during the composting of cattle manure

María Gómez-Brandón*, Cristina Lazcano, Jorge Domínguez

Departamento de Ecología y Biología Animal, Universidad de Vigo, Vigo E-36310, Spain

Received 27 February 2007; received in revised form 22 June 2007; accepted 26 June 2007

Available online 8 August 2007

Abstract

We examined chemical, microbiological and biochemical parameters in order to assess their effectiveness as stability and maturity indicators during the composting process of cattle manure. The composting material obtained after 15 d in trenches and at different times during the maturation phase (i.e. 80, 180 and 270 d) were analyzed. We found that the material collected at the end of the active phase was inadequate to be applied to soil as organic amendment due to its high content of NH_4^+ , its high level of phytotoxicity and the low degree of organic matter stability. After a maturation period of 80 d, the stability of the sample increased. This was shown by a reduction in the dissolved organic carbon (DOC) content and NH_4^+ concentration and also by a reduction in the microbial activity and biomass; however, 180 d of composting were not sufficient to reduce the phytotoxicity to levels consistent for a safe soil application. Among the various parameters studied, the change in DOC with composting time gave a good indication of stability.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: DOC; Mineral N; Microbial biomass; Basal respiration; Enzymatic activities; Phytotoxicity

1. Introduction

Cattle manure is a valuable resource as a soil fertilizer, providing a high content of macro- and micronutrients for crop growth, and represents a low-cost alternative to mineral fertilizers (Sharpley and Smith, 1995). However, the overproduction of organic wastes by cattle breeding has led to inappropriate disposal practices; for example, their indiscriminate application to agricultural fields and their improper timing of application, that is, they are not applied when it would be most beneficial for crops. These practices could cause serious environmental problems that could include an excessive input of potentially harmful trace metals, inorganic salts and pathogens (Hutchison et al., 2005); an increase in nutrient loss from soils through leaching, erosion and runoff due to not considering the nutrient requirements of crops (Vervoort et al., 1998);

and the emission of hydrogen sulphide, ammonia and other toxic gases (Salazar et al., 2005).

The composting process may significantly reduce the environmental problems associated with the management of manures by transforming them into a safer and more stabilized material for application to soil (Carr et al., 1995). To obtain high quality compost it is necessary to understand the changes that the material undergoes with the composting process. The stability and maturity of the compost are essential for its successful application, particularly for composts used in high value horticultural crops (Wang et al., 2004).

The terms stability and maturity are usually used interchangeably to describe the degree of decomposition and transformation of the organic matter in compost (Zmora-Nahum et al., 2005), despite the fact they describe different properties of the composting substrate. Stability is strongly related to the rate of microbial activity in compost, and is evaluated by different respirometric measurements (Lasaridi and Stentiford, 1998) and/or by studying the transformations in the chemical characteristics of compost organic

* Corresponding author. Tel.: +34 986 812593; fax: +34 986 812556.
E-mail address: mariagomez@uvigo.es (M. Gómez-Brandón).

matter (Pichler and Kögel-Knabner, 2000). Respirometric tests have been shown to be adequate for assessing compost stability because they are able to measure the extent of which readily biodegradable organic matter has decomposed during the composting process (Adani et al., 2004). Compost maturity generally refers to the degree of decomposition of phytotoxic organic substances produced during the active composting phase and to the absence of pathogens and viable weed seeds (Wu et al., 2000). Both these properties are critical for the quality and marketability of the final product.

The application of unstable compost to soil may produce a competition for oxygen between microbial biomass and plant roots/seeds. This fact can deprive plant roots/seeds of oxygen, and lead to the production of H_2S and NO_2^- (Mathur et al., 1993). Another problem is nitrogen starvation of plants as microorganisms scavenge soil N to make up for the deficit resulting from the application of unstable compost with a high C to N ratio. The phytotoxicity of unstable composts represents another major problem; this is due to the emission of ammonia and the presence of other phytotoxic substances like phenolic compounds and ethylene oxide that is synthesized during the decomposition of unstable compost in soil. Low-molecular weight organic acids (i.e. acetic, propionic and butyric acids) produced by the anaerobic digestion of the organic matrix are also responsible for compost phytotoxicity (Fuchs, 2002).

Management of the composting process must consider the potential agronomic value of the end product and its suitability for plant crops by evaluating its degree of maturity. Biological methods involving seed germination tests and plant growth bioassays have been used to evaluate the maturity of compost (Cooperband et al., 2003). This is a tedious work and there are disagreements regarding the ability of these tests to determine compost maturity (Brewer and Sullivan, 2003).

A large variety of techniques have been reported for the determination of compost stability (Wang et al., 2004). Chemical parameters such as pH, electrical conductivity (EC), cation exchange capacity, dissolved organic carbon (DOC) and the ratios of C to N and NH_4^+ to NO_3^- have been applied as indicators of stability. Since stabilization implies the formation of humic-like substances, humification indexes are generally accepted as a criterion of stability, but their absolute values vary greatly among composts of different source materials. Moreover, their determination requires proper separation of the non-humic fraction from the fulvic acid fraction because other compounds with similar structure to humic substances but different biological meaning (i.e. lignin residues, quinones, polypeptides, fats, etc.) can be extracted (Sánchez-Monedero et al., 1999). Stability indicators based on the study of microbial biomass and its activity have also been proposed. Mondini et al. (2006) reported that microbial biomass can be used as a stability parameter in ligno-cellulosic waste composts because it clearly reflects the transformation of organic

matter during the composting process. Respiration (CO_2 evolution rate and/or O_2 uptake rate) is a general measure of microbial activity, and it has been widely used to evaluate the stability of compost (Gómez et al., 2006). The ATP content and enzyme activities are also useful as indicators of compost stability (Tiquia et al., 2002; Boulter-Bitzer et al., 2006).

The use of different parameters appropriate to determine the maturity and/or stability of composts will allow us to broaden our knowledge about the composting process. Therefore, the two major objectives of this study were (a) to describe the chemical, microbiological and biochemical changes during the industrial composting of cattle manure and (b) to compare different parameters with respect to their ability to evaluate compost stability and maturity during the industrial composting of cattle manure.

2. Materials and methods

2.1. Source materials and composting process

This study followed the composting process of fresh cattle manure obtained from the agricultural cattle complex “Energía Viva, S.A.” in León, Spain. The researchers did not control the composting operation or attempt to influence the course of the composting process, which involved an active phase of 15 d, followed by a maturation stage in piles for 270 d.

Cattle manure subject to the active phase in five trenches with approximate dimensions of 42 m long, 1.8 m wide and 4.5 m high where each contained approximately 300 m^3 of material. Throughout the process, these trenches were aerated from the bottom with forced air through a blower in order to induce air convection movement into the material and deliver oxygen to microorganisms. The functioning of the air blower varied as a function of the temperature: (i) continuous aeration when the temperature of the composting mass overcame the value of 60°C ; (ii) intermittent aeration according to a preset cycle of 5 min aeration and 5 min pause when the temperature was found between 55°C and 60°C ; and (iii) intermittent aeration according to a preset cycle of 5 min aeration followed by 10 min pause when the temperature was below 55°C . The forced ventilation was combined with daily turnings in order to homogenize the composting mass and to avoid the substrate compaction and the subsequent low porosity and deficient air distribution. The composting material was watered with water and the moisture content was controlled daily and kept within the range of recommended values (45–65%; Miller, 1993).

During the curing phase, the composting mixture from each trench was piled up and left to mature in maturation piles (50 m long, 2 m wide and 2 m high) up to 270 d in a space covered on top by a ceiling with the sides opened. These piles were turned for aeration twice a month and sporadically watered with leachates from the cattle farm. Samples were collected at 10 random locations at 15, 80,

180 and 270 d and thoroughly mixed to generate composite samples. All the samples were stored in sealed plastic containers that were kept at 5 °C until they were analyzed. The initial fresh cattle manure was also analyzed for comparison.

2.2. Chemical analyses

EC and pH were analyzed in water extracts (1:10, w/v). Total C and N contents were analyzed on a Carlo Erba 1500C/N analyzer on dried samples. DOC was determined in 0.5 M K₂SO₄ extracts (1:50, w/v) by heat digestion (150 °C, 30 min) with sulphuric acid and potassium dichromate and read in a Bio-Rad Microplate Reader 550 at 590 nm. Inorganic nitrogen (NH₄⁺ and NO₃⁻) was determined in 0.5 M K₂SO₄ extracts (1:5, w/v) using the modified indophenol blue technique (Sims et al., 1995) with a Bio-Rad Microplate Reader 550. The content of P was analyzed in ammonium bicarbonate-diethylenetriamine-pentaacetic acid (AB-DTPA) extracts (1:6, w/v) by atomic absorption spectrophotometry (Soltanpour and Schwab, 1977).

2.3. Microbiological and biochemical analyses

Microbial biomass carbon (C_{mic}) was determined by the chloroform fumigation–extraction method (Vance et al., 1987) on moist samples (5 g fresh weight). The filtered extracts of both fumigated and non-fumigated samples were analyzed for soluble organic C using a Microplate Reader (Bio-Rad Microplate Reader 550, 590 nm). C_{mic} was estimated as the difference between the organic C extracted from the fumigated and the non-fumigated sample, multiplied by the K₂SO₄ extraction efficiency factor for microbial C ($K_c = 2.64$).

Ergosterol is a membrane-bound molecule commonly used as a fungal biomarker (Bååth and Anderson, 2003). The ergosterol content of the samples was extracted by the microwave assisted extraction method and determined by HPLC analysis (Young, 1995).

Microbial activity was assessed by measuring the rate of CO₂ evolution from the samples (5 g fresh weight) after 6 h of incubation. The evolved CO₂ was trapped in 0.02 M NaOH and then measured by titration with HCl to a phenolphthalein endpoint, after adding excess BaCl₂ (Anderson, 1982).

Alkaline phosphomonoesterase activity was estimated by determining the *p*-nitrophenol (PNP) released, after incubating the samples (1 g fresh weight) with *p*-nitrophenyl phosphate (0.025 M) for 1 h at 37 °C, in a Bio-Rad Microplate Reader 550 at 400 nm (Eivazi and Tabatabai, 1972). β-glucosidase activity was assessed by determining the PNP released, after incubating the samples (1 g fresh weight) with β-D-glucopyranoside (0.025 M) for 1 h at 37 °C, in a Bio-Rad Microplate Reader 550 at 400 nm (Eivazi and Tabatabai, 1988). Protease activity was measured by determining the amino acids released, after incubating the sam-

ples (1 g fresh weight) with sodium caseinate (2%) for 2 h at 50 °C, using Folin–Ciocalteu reagent, in a Bio-Rad Microplate Reader 550 at 700 nm (Ladd and Butler, 1972). Cellulase activity was estimated by determining the reducing sugars released after incubating the samples (5 g fresh weight) with carboxymethyl cellulose sodium salt (0.7%) for 24 h at 50 °C, in a Bio-Rad Microplate Reader at 690 nm (Schinner and Von Mersi, 1990).

2.4. Phytotoxicity assay

The phytotoxicity of the samples was determined in distilled water extracts (1:5, w/v) following the method of Zucconi and de Bertoldi (1987). The extracts were agitated vigorously for 1 h and then centrifuged for 15 min at 10000 rpm in order to separate the phases. The supernatant was collected and filtered through a 0.45 μm membrane filter. The extracts were diluted to 30% and used as germination media. One milliliter of the germination solution was pipetted into a sterilized Petri dish lined with Whatman #1 filter paper. Ten seeds of garden cress (*Lepidium sativum* L.), which is one of the most sensitive test species for evaluating the phytotoxicity (Gehring et al., 2003), were evenly distributed on the filter paper and incubated 24 h at 20–25 °C in the dark. Then, the germination was stopped by adding 1 ml of ethanol. The seed germination percentage and root elongation of *L. sativum* were also measured in distilled water and used as control. Finally, the germination index (GI), expressed as percentage of control, was calculated based on relative seed germination percentage and relative root elongation.

2.5. Statistical analyses

The statistical analysis of data was carried out using the SPSS 11.0 program for Windows. A normality test was made for all the parameters prior to analyzing the variance. The results were submitted to an ANOVA test in order to determine changes in the variables with the composting time. A Tukey's test was used for testing significant statistical differences among composting times. The relationships between variables were defined by regression analysis. A principal components analysis was carried out to summarize the results obtained with the chemical, biochemical and microbiological parameters. Two principal components (PC1 and PC2) were used for this analysis.

3. Results and discussion

3.1. Chemical changes

The main chemical properties of the initial cattle manure and the composting mixture after 15, 80, 180 and 270 d are shown in Table 1. An increase in the pH values was recorded during the active phase, suggesting the alkalization of the manure as a consequence of the release of ammonia from the degradation and mineralization of

Table 1
Variation in the chemical properties of cattle manure with composting time

	Composting time (days)				
	Thermophilic		Maturation		
	0	15	80	180	270
EC (mS cm ⁻¹)	1.30 ± 0.08c	2.10 ± 0.11b	2.80 ± 0.23a	2.90 ± 0.16a	3.00 ± 0.12a
pH	8.26–8.30b	8.07–8.86b	9.20–9.40a	9.50–9.75a	9.48–9.72a
Total C (%)	39.90 ± 0.28a	38.50 ± 0.26a	32.15 ± 0.53b	33 ± 0.53b	25 ± 1.22c
Total N (%)	2.40 ± 0.09b	2.20 ± 0.03b	2.90 ± 0.05a	3.10 ± 0.04a	2.40 ± 0.13b
C to N ratio	17 ± 0.74a	17.50 ± 0.33a	11.40 ± 0.34b	10.75 ± 0.09b	10.60 ± 0.12b
DOC (mg kg ⁻¹ dw)	7000 ± 700a	7300 ± 280a	3400 ± 610b	3100 ± 250b	2200 ± 10b
NH ₄ ⁺ (mg kg ⁻¹ w)	600 ± 90b	1200 ± 10a	300 ± 40c	120 ± 10c	250 ± 30c
NO ₃ ⁻ (mg kg ⁻¹ dw)	20 ± 10b	700 ± 60a	20 ± 10b	50 ± 10b	70 ± 10b
NH ₄ ⁺ to NO ₃ ⁻ ratio	30 ± 14a	1.71 ± 0.58b	15 ± 7.10a	2.4 ± 0.96b	3.57 ± 0.89b
Available P (mg kg ⁻¹)	170 ± 10b	350 ± 20a	120 ± 30bc	110 ± 10bc	70 ± 10c

Values are means ± standard error ($n = 5$). Values within the same row followed by the same letter are not significantly different according to Tukey's test ($P = 0.05$).

organic compounds. During the maturation stage, pH reached higher levels than in the initial cattle manure and the sample collected at the end of the active phase. This increase is not typical in the maturation phase, where it is expected that the pH drops to neutral values and then stabilizes. The watering of the maturation piles with the leachates from the cattle farm, and the turnings of these piles could be responsible of these results. The pH values reached during the maturation are rather high and this fact could have important implications on the fertility and productivity of soils subjected to compost amendment, as well as on the development of pH-sensitive plants (Boulter-Bitzer et al., 2000). Moreover, acceptable pH ranges should be within tolerable levels to microorganisms, i.e. bacteria generally need a pH range of 6–7.5; fungi can tolerate a wider range, 5.5–8.0; and actinomycetes 5.0–9.0. Our composting material collected at different times during the maturation phase presented pH ranges that exceeded the previous ones; thus, the addition of these materials to soil may also strongly influence the soil microflora.

The EC also affects the quality of composts in a large way because it reflects their salinity and suitability for crop growth. This parameter increased after the active phase probably due to the release of soluble salts like ammonium and phosphate resulting from the decomposition of easily biodegradable organic substrates. As in the case of the pH, higher levels of EC were found during the maturation stage. The addition of liquid animal wastes to the maturation piles and the turning of these piles could have been responsible for the increase in EC. In spite of this increase, the EC of compost obtained after 270 d of maturation did not exceed the limit value of 3 mS cm⁻¹ indicating a material that could be safely applied to soil (Soumaré et al., 2002).

When an organic waste is composted, the C to N ratio generally decreases throughout the composting process due to the C losses as CO₂ and then stabilizes in the range of 10–15 (Chefetz et al., 1996). As the total N content changed slightly with the composting time, in the present study, a significant reduction of this ratio during the maturation phase was mainly due to the depletion of easily degradable

carbon compounds as outlined by DOC dynamics (Table 1). The rate of decrease in DOC concentration depends on the source material and the composting technique utilized. The amount of DOC during the composting process is related to the equilibrium between various reactions which increase or decrease its concentration. The degradation of solid polymeric material in the composting substrate may lead to the formation of soluble organic matter, which would increase the DOC concentration. On the other hand, the reduction in DOC depends on the continuous mineralization of soluble organic compounds, and the repolymerization and condensation pathways that lead to the formation of complex organic substrates with low solubility in water which tend to flocculate out the solution (Said-Pullicino and Gigliotti, 2007).

Mineral N (NH₄⁺ and NO₃⁻) increased significantly during the active phase which indicates an intense mineralization. On the contrary, at maturation stage, the concentration of NH₄⁺ greatly decreased due to its volatilisation as NH₃ as a result of the high pH observed during this phase and most likely because of the frequent turning. Low levels of NO₃⁻ were also found during the maturation phase. This fact may have been the result of leaching of NO₃⁻ with the watering of the maturation piles. Brewer and Sullivan (2003) reported a decrease in NO₃⁻ in yard trimmings compost obtained after 133 d of maturation probably due to the leaching from saturated compost or denitrification. Compared to the initial cattle manure, the NH₄⁺ to NO₃⁻ ratio greatly decreased during the active phase due to the high levels of NO₃⁻ detected in this stage. After 80 d of maturation, this ratio increased with respect to the sample collected at the end of the active phase. However, lower values of this ratio were reported in the samples collected after 180 and 270 d of maturation mainly due to the low content of NH₄⁺.

3.2. Microbiological and biochemical changes

The decreasing trend of microbial biomass throughout the composting process (Fig. 1a) was in agreement with




Fig. 1. Changes in (a) Cmic, (b) ergosterol content and (c) basal respiration of the initial cattle manure ($t = 0$) and the composting material collected at the end of the active phase and at different times during the maturation stage (i.e. 80, 180 and 270 d). Values are means \pm standard error ($n = 5$). Different letters indicate significant differences at $P < 0.05$ (Tukey HSD).

other works (García et al., 1992; Insam et al., 1996; Klamer and Bååth, 1998), reporting results obtained with different biomass quantification methods (fumigation–extraction, substrate-induced respiration, ATP content, total phospholipid fatty acids content). The fungal biomass measured as ergosterol content also decreased during the active phase and maturation stage compared to the initial cattle manure (Fig. 1b).

The microbial activity measured as basal respiration reached a maximum value of $5000 \text{ mg CO}_2 \text{ kg}^{-1}$ organic matter during the active phase, which could be attributed to the presence of easily degradable materials that stimulate the microbial community of the initial cattle manure. Moreover, the release of labile compounds resulting from the oxidative biodegradation of the matrix during the composting process could also explain the higher value obtained on day 15 with respect to the initial feedstock (Said-Pullicino et al., 2007). However, lower rates of respiration indicative of minor microbial activity were recorded

after the maturation period of 80 d (Fig. 1c). The decrease of microbial activity was corroborated by the reduction in the microbial biomass as indicated by the low levels of Cmic during the maturation period.

Enzymatic parameters also reflect the activity of the microbial community and indicate the ability of composting to degrade a wide range of common organic substrates (Mondini et al., 2004). Important enzymes involved in the composting process include cellulases, which depolymerise cellulose; β -glucosidases which hydrolyse glucosides; amidohydrolases, proteases and ureases involved in N mineralization; and phosphomonoesterases and arylsulphatases that remove phosphate and sulphate groups from organic compounds.

In our study, the alkaline phosphomonoesterase activity greatly decreased during the active phase probably due to the feedback inhibition of this enzyme by inorganic phosphate (Ayuso et al., 1996). The high levels of extractable P found in the sample collected at the end of the active phase support this hypothesis (Table 1). However, compared to this sample, this enzyme activity was significantly higher after 180 and 270 d of maturation (Fig. 2a). This could be due to the decrease in extractable P content during this period, as well as due to the formation of an enzyme–humus complex, which would make this enzyme more resistant to denaturation (Mondini et al., 2004). Protease activity decreased throughout the process (Fig. 2b), which is indicative of a reduction in the substrate for protease capable of activating the synthesis of this enzyme (Diaz Burgos and Polo, 1991). Compared to the initial cattle manure, lower levels of cellulase activity were reported during the active and maturation stages (Fig. 2c). On the other hand there were no significant differences in β -glucosidase activity with the composting time (Fig. 2d). It is assumed that soil cellulases and β -glucosidases are mainly produced by fungi (Hayano, 1986). In the present study, a reduction in the fungal biomarker ergosterol was observed during the active phase and the maturation stage (Fig. 1b). We found a significant correlation between ergosterol and cellulase activity ($R^2 = 0.50$, $P = 0.0001$), but not with the β -glucosidase activity ($R^2 = 0.05$, $P = 0.164$). One possible explanation is the protection of β -glucosidase through the formation of complexes with the humic substances, which made the enzyme more resistant to physical and microbial degradation.

3.3. Summary of chemical, microbiological and biochemical changes using a principal components analysis

PC1 explained 60% of the variance and PC2 22% for a total explained variance of 82%. The variables responsible for the changes along PC1 included pH, EC, total C and N, C to N ratio, DOC, NH_4^+ , NO_3^- , NH_4^+ to NO_3^- ratio, extractable P, Cmic, basal respiration and protease activity. All of them were positively correlated with this function, except pH, EC and total N. The variables responsible for the changes along PC2 included the




Fig. 2. Changes in (a) alkaline phosphomonoesterase, (b) protease, (c) cellulase and (d) β -glucosidase activities of the initial cattle manure ($t = 0$) and the composting material collected at the end of the active phase and at different times during the maturation stage (i.e. 80, 180 and 270 d). Values are means \pm standard error ($n = 5$). Different letters indicate significant differences at $P < 0.05$ (Tukey HSD).

phosphomonoesterase and cellulase activities and ergosterol, which were positively correlated with this function. The sample collected at the end of the active phase separated from the initial cattle manure along PC2 (Fig. 3). This change was highlighted by a decrease in the concentration of ergosterol and phosphomonoesterase and cellulase activities during the active phase. However, the composting




Fig. 3. Principal components analysis of the chemical, microbiological and biochemical parameters of the initial cattle manure ($t = 0$) and the composting material collected at the end of the active phase and at different times during the maturation stage (i.e. 80, 180 and 270 d).

material collected at different times during the maturation phase separated from the initial cattle manure and the sample collected at the end of the active phase mainly along PC1. In this case, the change was characterized by an increase in the pH, EC and total N and a decrease in the total C, C to N ratio, DOC, NH_4^+ , NO_3^- , extractable P, Cmic, basal respiration and protease activity during the maturation period.

3.4. Phytotoxicity bioassay

The germination index was 0% in the initial cattle manure, and reached a value of 24% during the active phase. Then, at maturation stage, a GI of 87% was recorded after 270 d of maturation (Fig. 4). Thus, more than 180 d were needed to overcome the threshold limit of 60% stated by Zucconi and de Bertoldi (1987) to reduce the phytotoxicity to levels consistent for a safe soil application.

3.5. Evaluation of stability and maturity parameters

The greatest concern regarding the evaluation of maturity is that this parameter does not only depend on the type of material and composting process utilized but also on the target use of the final product. As the composting of animal manures increases, due to its practical characteristic as a method of recycling, it becomes more and more important to determine the appropriate use of each method for the evaluation of the stability and maturity of the compost.

The application of chemical parameters to assess the stability of compost is a common practise in the research on




Fig. 4. Changes in the germination index of the initial cattle manure ($t = 0$) and the composting material collected at the end of the active phase and at different times during the maturation stage (i.e. 80, 180 and 270 d). Values are means \pm standard error ($n = 5$). Different letters indicate significant differences at $P < 0.05$ (Tukey HSD).

composting. To be a good indicator, a chemical parameter should fulfil the following requirements: (a) follow a consistent trend during the composting process; (b) provide reference, critical or threshold values; (c) require relatively rapid and cheap analytical procedures and (d) be easily interpretable. Some of the parameters frequently referred to in the literature regarding the evaluation of compost stability are discussed in this study.

C to N ratio may not be a good indicator of compost stability because it can level off before the compost stabilizes (Namkoong et al., 1999). For example, when wastes rich in nitrogen are used as source material for composting, like sewage sludges or manures, the C to N ratio can be within the values of a stable compost even though it may still be unstable. Zmora-Nahum et al. (2005) reported a C to N ratio lower than the cutoff value of 15 very early during the composting of cattle manure, while important stabilization processes were still taking place.

DOC generally contains organic compounds having different susceptibilities to microbial degradation and different phytotoxic properties. For this reason DOC composition may have an important role in determining the stabilization process. In the present study, DOC concentrations during the maturation phase were lower than the threshold value of 4000 mg kg^{-1} suggested by Zmora-Nahum et al. (2005) for a stable compost showing that the composting material was stabilized after 80 d from the point of view of DOC content (Table 1). In addition, we found a significant correlation between DOC and respiration ($R^2 = 0.60$, $P = 0.0001$). Chica et al. (2003) also showed DOC to be highly correlated to respiration.

Compost stability can be determined in terms of nitrification, which mainly takes place during the maturation stage when temperatures are close to the ambient. At the end of the composting process the content of NO_3^- should be higher than that of NH_4^+ , indicating that the process has been performed under adequate aeration conditions (Bernal et al., 1998). Our composting material collected at different times during the maturation phase did not exceed the limit value of 400 mg kg^{-1} suggested by Zucconi and de Bertoldi (1987) for the concentration of NH_4^+ in stable composts; but, the levels of NO_3^- were lower than expected during the maturation phase (Table 1). The NH_4^+ to NO_3^- ratio has also been used to estimate the compost stability (Bernal et al., 1998), giving a limit value of 0.16 for stable composts. In our case, this value was exceeded during the maturation phase due to the low content of NO_3^- detected during this stage (Table 1). As mineral nitrogen forms changed irregularly with the composting time, they cannot be reliable indicators.

After a maturation period of 80 d, similarly to DOC, the stability assessed by the rate of microbial respiration increased, because lower rates of respiration were reported in the composting material collected at different times during the maturation stage (Fig. 1c). Wang et al. (2004) also reported low rates of respiration, indicative of highly stabilized composts, towards the end of the composting of dairy and pig manures. Adani et al. (2004) proposed a dynamic respiration index as an accurate indicator to measure the stability of composts resulting from different starting materials. They reported threshold values of 1000 and $500 \text{ mg O}_2 \text{ kg}^{-1} \text{ organic matter h}^{-1}$ to indicate medium and high stability, respectively. Other studies have established that the measurements of microbial respiration can be problematic as they are very sensitive to changes of moisture, temperature, and oxygen and nitrogen availability (Herrmann and Shann, 1993).

Wu et al. (2000) reported that the low CO_2 evolution is not always an indicator of a non-phytotoxic compost. Therefore, the evaluation of compost stability based on CO_2 evolution, and the maturity based on seed germination, are two different parameters of compost quality. In our study, despite of the fact that low rates of respiration, indicative of highly stabilized materials, were reported after 80 d of maturation, more than 180 d were needed to overcome the threshold limit of 60% stated by Zucconi and de Bertoldi (1987) for a compost to be considered phytotoxin-free (Fig. 4). The source material and the composting condition as well as the watering of the maturation piles with leachates from the cattle farm could have required more time to break down the phytotoxic substances.

Considering the key roles of microorganisms in the composting process, the use of microbiological properties as stability indicators is not surprising. Enzymatic activities play an important role during the composting process, as they are implicated in the biological and biochemical processes through which the initial organic substrates are transformed into the end product (Tiquia, 2002). As a

consequence, specific enzymatic activities could provide a way of characterizing the composting process with relation to the rate of transformation of organic residues and the stability of the end product. Mondini et al. (2004) reported that the change in the location of enzymes throughout the composting process, i.e. from extracellular to complexed with humic-like substances, might be useful at the moment of evaluating compost stability, taking into account that not all enzymes will be as equally reliable as a stability index. Despite of the fact that the measurement of enzymatic activities is easy, quick and inexpensive, it is difficult to establish general threshold values to apply enzymatic activities as indicators of compost stability due to the widely different organic substrates involved in the composting processes. Thus, for compost characterization, it is necessary to follow the dynamics of enzymatic activities over time.

Having discussed the previous parameters, determining the change in DOC content with composting time seems to be the most suitable measurement to evaluate the stability of the composting material. It fulfilled the largest number of requirements, it followed a consistent trend and reached a critical value; moreover, it was neither time-consuming nor expensive and was easy to interpret. However, this does not mean that this measurement is equally accurate to evaluate the stability of all source materials and full composting facilities. The creation of databases, showing which protocols are most effective taking into account the source material, the composting conditions and if the experiment was done in a laboratory or at full scale will help us to choose correctly the different parameters for the evaluation of compost quality.

4. Conclusions

The active phase was accompanied by a significant increase in EC, mineral N, extractable P and basal respiration. However, lower levels of ergosterol and phosphomonoesterase, protease and cellulase activities were found during this phase. The maturation period was characterized by an increase in pH, EC and phosphomonoesterase activity and a decrease in C to N ratio, DOC, mineral N, extractable P, basal respiration and protease activity. The turning of maturation piles and the sporadic addition of leachates from the cattle farm to these piles influenced several parameters such as the pH and NO_3^- content resulting in values that are not typical from the maturation stage. Moreover, a maturation phase of 80 d was enough to obtain a stable compost as outlined by several parameters, but not to obtain a mature compost (i.e. with a low degree of phytotoxicity).

Acknowledgements

María Gómez Brandón is financially supported by a FPU fellowship from Ministerio de Educación. The authors thank the personnel of “Energía Viva, S.A.” for

having let us follow the composting process in their facility. The authors thank Paul Fraiz for having revised the English in this article. The authors also thank the editor and the two anonymous reviewers for helping us to improve the quality of this work.

References

- Adani, F., Gonfalonieri, R., Tambone, F., 2004. Dynamic respiration index as a descriptor of the biological stability of organic wastes. *J. Environ. Qual.* 33, 1866–1876.
- Anderson, J.P.E., 1982. Soil respiration. In: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.), *Methods of Soil Analysis, Part 2. Chemical and Biological Properties*. American Society of Agronomy and Soil Science Society of America, Madison, WI, pp. 841–845.
- Ayuso, M., Pascual, J.A., García, C., Hernández, T., 1996. Evaluation of urban wastes for agricultural use. *Soil Sci. Plant Nutr.* 42, 105–111.
- Bååth, E., Anderson, T.H., 2003. Comparison of soil fungal/bacterial ratios in a pH gradient using physiological and PLFA- techniques. *Soil Biol. Biochem.* 35, 955–963.
- Bernal, M.P., Paredes, C., Sánchez-Monedero, M.A., Cegarra, J., 1998. Maturity and stability parameters of composts prepared with a wide range of organic wastes. *Bioresour. Technol.* 63, 91–99.
- Boulter-Bitzer, J.I., Boland, G.J., Trevors, J.T., 2000. Compost: a study of the development process and end-product potential for suppression of turfgrass disease. *World J. Microbiol. Biotechnol.* 16, 115–134.
- Boulter-Bitzer, J.I., Trevors, J.T., Boland, G.J., 2006. A polyphasic approach for assessing maturity and stability in compost intended for suppression of plant pathogens. *Appl. Soil Ecol.* 34, 65–81.
- Brewer, L.J., Sullivan, D.M., 2003. Maturity and stability evaluation of composted yard trimmings. *Compost Sci. Util.* 11, 96–112.
- Carr, L., Grover, R., Smith, B., Richard, T., Halbach, T., 1995. Commercial and on-farm production and marketing of animal waste compost products. In: Steele, K. (Ed.), *Animal Waste and the Land–Water Interface*. Lewis Publishers, Boca Raton, pp. 485–492.
- Chefetz, B., Hatcher, P.G., Hadr, Y., Chen, Y., 1996. Chemical and biological characterization of organic matter during composting of municipal solid waste. *J. Environ. Qual.* 25, 776–785.
- Chica, A., Mohedo, J.J., Martín, M.A., Martín, A., 2003. Determination of the stability of MSW compost using a respirometric technique. *Compost Sci. Util.* 11, 169–175.
- Cooperband, L.R., Stone, A.G., Fryda, M.R., Ravet, J.L., 2003. Relating compost measures of stability and maturity to plant growth. *Compost Sci. Util.* 11, 113–124.
- Díaz Burgos, M.A., Polo, A., 1991. Variaciones en la fracción orgánica durante el compostaje de lodos de depuradora. *Suelo y Planta* 1, 453–466.
- Eivazi, F., Tabatabai, M.A., 1972. Phosphatases in soil. *Soil Biol. Biochem.* 9, 167–172.
- Eivazi, F., Tabatabai, M.A., 1988. Glucosidases and galactosidases in soil. *Soil Biol. Biochem.* 20, 601–606.
- Fuchs, J.G., 2002. Practical use of quality compost for plant health and viability improvement. In: Insam, H., Riddech, N., Klammer, S. (Eds.), *Microbiology of Composting*. Springer Verlag, Heidelberg, pp. 435–444.
- García, C., Hernández, T., Costa, C., Ayuso, M., 1992. Evaluation of the maturity of municipal waste compost using simple chemical parameters. *Commun. Soil Sci. Plant Anal.* 23, 1501–1502.
- Gehring, M.M., Kewada, V., Coates, N., Downing, T.G., 2003. The use of *Lepidium sativum* in a plant bioassay system for the detection of microcystin-LR. *Toxicon* 41, 871–876.
- Gómez, R., Vázquez-Lima, F., Sánchez-Ferrer, A., 2006. The use of respiration indices in the composting process: a review. *Waste Manage. Res.* 24, 37–47.
- Hayano, K., 1986. Cellulase complex in tomato field soil: introduction, localization and some properties. *Soil Biol. Biochem.* 18, 215–219.

- Herrmann, R.F., Shann, J.R., 1993. Enzyme activities as indicators of municipal solid waste compost maturity. *Compost Sci. Util.* 4, 54–63.
- Hutchison, M.L., Walters, L.D., Avery, S.M., Munro, F., Moore, A., 2005. Analyses of livestock production, waste storage, and pathogen levels and prevalences in farm manures. *Appl. Environ. Microbiol.* 71, 1231–1236.
- Insam, H., Amor, K., Renner, M., Crepaz, C., 1996. Changes in functional abilities of the microbial community during composting of manure. *Microbial Ecol.* 31, 77–87.
- Klamer, M., Bååth, E., 1998. Microbial community dynamics during composting of straw material studied using phospholipid fatty acid analysis. *FEMS Microbiol. Ecol.* 27, 9–20.
- Ladd, J.N., Butler, J.H.A., 1972. Short-term assays of soil proteolytic enzymes activities using proteins and dipeptide derivatives as substrates. *Soil Biol. Biochem.* 4, 19–30.
- Lasaridi, K.E., Stentiford, E.I., 1998. A simple respirometric technique for assessing compost stability. *Water Res.* 32, 3717–3723.
- Mathur, S.P., Owen, G., Diné, H., Schnitzer, M., 1993. Determination of compost biomaturity. *Biol. Agric. Hortic.* 10, 65–85.
- Miller, F.C., 1993. Composting as a process based on the control of ecologically selective factors. In: Blaine, F., Metting, J. (Eds.), *Soil Microbial Ecology – Applications in Agricultural and Environmental Management*. Marcel Dekker, New York, pp. 515–543.
- Mondini, C., Fornasier, F., Sinicco, T., 2004. Enzymatic activity as a parameter for the characterization of the composting process. *Soil Biol. Biochem.* 36, 1587–1594.
- Mondini, C., Sánchez-Monedero, M.A., Sinicco, T., Leita, L., 2006. Evaluation of extracted organic carbon and microbial biomass as stability parameters in ligno-cellulosic waste composts. *J. Environ. Qual.* 35, 2313–2320.
- Namkoong, W., Hwang, E.Y., Cheong, J.G., Choi, J.Y., 1999. A comparative evaluation of maturity parameters of food waste composting. *Compost Sci. Util.* 7, 55–62.
- Pichler, M., Kögel-Knabner, I., 2000. Chemolytic analysis of organic matter during aerobic and anaerobic treatment of municipal solid waste. *Waste Manage.* 29, 1337–1344.
- Said-Pullicino, D., Gigliotti, G., 2007. Oxidative biodegradation of dissolved organic matter during composting. *Chemosphere* 68, 1030–1040.
- Said-Pullicino, D., Erriquens, F.G., Gigliotti, G., 2007. Changes in the chemical characteristics of water-extractable organic matter during composting and their influence on compost stability and maturity. *Bioresour. Technol.* 98, 1822–1831.
- Salazar, F.J., Chadwick, D., Pain, B.F., Hatch, D., Owen, E., 2005. Nitrogen budgets for three cropping systems fertilised with cattle manure. *Bioresour. Technol.* 96, 235–245.
- Sánchez-Monedero, M.A., Roig, A., Cegarra, J., Bernal, M.P., 1999. Relationships between water-soluble carbohydrate and phenol fractions and the humification indices of different organic wastes during composting. *Bioresour. Technol.* 78, 301–308.
- Schinner, F., Von Mersi, W., 1990. Xylanase-, CM-cellulase- and invertase activity in soil: an improved method. *Soil Biol. Biochem.* 22, 511–515.
- Sharpley, A.M., Smith, S.J., 1995. Nitrogen and phosphorus in soils receiving manure. *Soil Sci.* 159, 253–258.
- Sims, G.K., Ellsworth, T.R., Mulvaney, R.L., 1995. Microscale determination of inorganic nitrogen in water and soil extracts. *Commun. Soil Sci. Plant Anal.* 26, 303–316.
- Soltanpour, P.N., Schwab, P.A., 1977. A new soil test for simultaneous extraction of macro- and micronutrients in alkaline soils. *Commun. Soil Sci. Plant Anal.* 8, 195–207.
- Soumaré, M., Demeyer, A., Tack, F.M.G., Verloo, M.G., 2002. Chemical characteristics of Malian and Belgian solid waste composts. *Bioresour. Technol.* 81, 97–101.
- Tiquia, S.M., 2002. Evolution of extracellular enzyme activities during manure composting. *J. Appl. Microbiol.* 92, 764–775.
- Tiquia, S.M., Wan, J.H.C., Tam, N.F.Y., 2002. Dynamics of yard trimmings composting as determined by dehydrogenase activity, ATP content, arginine ammonification, and nitrification potential. *Process Biochem.* 37, 1057–1065.
- Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil microbial biomass C. *Soil Biol. Biochem.* 19, 703–707.
- Vervoort, R.V., Radcliffe, D.E., Cabrera, M.L., Latimore Jr., M., 1998. Nutrient losses in surface and subsurface flow from pasture applied poultry litter and composted poultry litter. *Nutr. Cycl. Agroecosys.* 50, 287–290.
- Wang, P., Changa, C.M., Watson, M.E., Dick, W.A., Chen, Y., Hoitink, H.A.J., 2004. Maturity indices for composted dairy and pig manures. *Soil Biol. Biochem.* 36, 767–776.
- Wu, L., Ma, L.Q., Martinez, G.A., 2000. Comparison of methods for evaluating stability and maturity of biosolids compost. *J. Environ. Qual.* 29, 424–429.
- Young, J.C., 1995. Microwave-assisted extraction of the fungal metabolite ergosterol and total fatty acids. *J. Agric. Food Chem.* 43, 2904–2910.
- Zmora-Nahum, S., Markovitch, O., Tarchitzky, J., Chen, Y., 2005. Dissolved organic carbon (DOC) as a parameter of compost maturity. *Soil Biol. Biochem.* 37, 2109–2116.
- Zuconi, F., de Bertoldi, M., 1987. Compost specification for the production and characterization of compost from municipal solid waste. In: de Bertoldi, M., Ferranti, M.P., Hermite, P.L., Zuconi, F. (Eds.), *Compost: Production, Quality and Use*. Elsevier Applied Science Publishers, Barking, pp. 30–50.